Lecture 11 – Probability

DSC 10, Fall 2025

Disc today Ouiz 2 Weds

Agenda

We'll cover the basics of probability theory. This is a math lesson; take written notes .

Probability resources

Probability is a tricky subject. If it doesn't click during lecture or on the assignments, take a look at the following resources:

- Computational and Inferential Thinking, Chapter 9.5.
- Theory Meets Data, Chapters 1 and 2.
- Khan Academy's unit on Probability.

Probability theory

- Some things in life seem random.
 - e.g., flipping a coin or rolling a die 🐼.
- The **probability** of seeing "heads" when flipping a fair coin is $\frac{1}{2}$.
- One interpretation of probability says that if we flipped a coin infinitely many times, then $\frac{1}{2}$ of the outcomes would be heads.

Terminology

- **Experiment**: A process or action whose result is random.
 - e.g., rolling a die.
 - e.g., flipping a coin twice.
- Outcome: The result of an experiment.
 - e.g., the possible outcomes of rolling a six-sided die are 1, 2, 3, 4, 5,
 and 6.
 - e.g., the possible outcomes of flipping a coin twice are HH, HT, TH and TT.
- Event: A set of outcomes.
 - e.g., the event that the die lands on a even number is the set of outcomes {2, 4, 6}.
 - e.g., the event that the die lands on a 5 is the set of outcomes {5}.
 - e.g., the event that there is at least 1 head in 2 flips is the set of outcomes {HH, HT, TH}.

Terminology


- **Probability**: A number between 0 and 1 (equivalently, between 0% and 100%) that describes the likelihood of an event.
 - 0: The event never happens.
 - 1: The event always happens.
- $\bullet\,$ Notation: If A is an event, P(A) is the probability of that event.


Equally-likely outcomes

- P(even #) = 3 1
- ullet If all of the possible outcomes are equally likely, then the probability of A is

$$P(A) = rac{\# ext{ of outcomes satisfying } A}{ ext{total } \# ext{ of outcomes}}$$

• Example 1: Suppose we flip a fair coin 3 times. What is the probability we was





Conditional probabilities

- ullet Two events A and B can both happen. Suppose that we know A has happened, but we don't know if B has.
- If all outcomes are equally likely, then the conditional probability of B given A is:

$$A$$
 is:


$$P(B \text{ given } A) = \frac{\# \text{ of outcomes}}{\# \text{ of outcomes}}$$

$$\frac{\text{satisfying both } A \text{ and } B}{\text{satisfying } A}$$

• Intuitively, this is similar to the definition of the regular probability of B:

$$P(B) = \frac{\# \text{ of outcomes satisfying } B}{\text{total } \# \text{ of outcomes}}$$

if you restrict the set of possible outcomes to just those in event A. < P(BginnA)





Concept Check ✓ – Answer at <u>cc.dsc10.com</u>

$$P(B \text{ given } A) = rac{\# \text{ of outcomes}}{\# \text{ of outcomes}} rac{\text{satisfying both } A \text{ and } B}{\text{satisfying } A}$$

I roll a six-sided die and don't tell you what the result is, but I tell you that it is 3 or less. What is the probability that the result is even?

- A) $\frac{1}{2}$
- B) $\frac{1}{3}$
- C) $\frac{1}{4}$
- D) None of the above.






Probability that two events both happen

ullet Suppose again that A and B are two events, and that all outcomes are equally likely. Then, the probability that both A and B occur is

$$P(A \text{ and } B) = \frac{\# \text{ of outcomes}}{\text{total } \# \text{ of outcomes}}$$

• **Example 2**: I roll a fair six-sided die. What is the probability that the roll is 3 or less **and** even?




The multiplication rule


ullet The multiplication rule specifies how to compute the probability of both A and B happening, even if all outcomes are not equally likely.

$$P(A \text{ and } B) = P(A) \cdot P(B \text{ given } A)$$

• **Example 2, again**: I roll a fair six-sided die. What is the probability that the roll is 3 or less and even?




What if A isn't affected by B? $\stackrel{(\mathcal{G})}{=}$




$$P(A ext{ and } B) = P(A) \cdot P(B ext{ given } A)$$

- ullet What if knowing that A happens doesn't tell you anything about the likelihood of B happening?
 - Suppose we flip a fair coin three times.
 - The probability that the second flip is heads doesn't depend on the result of the first flip.








Independent events

equivalently if

ndependent: P(Bgivent) ullet Two events A and B are independent if $P(B ext{ given } A) = P(B)$, or

$$P(A \text{ and } B) = P(A) \cdot P(B)$$

Example 3: Suppose we have a coin that is biased, and flips heads with probability 0.7. Each flip is independent of all other flips. We flip it 5 times. What's the probability we see 5 heads in a row?



Probability that an event *doesn't* happen

- ullet The probability that A doesn't happen is 1-P(A) .
- For example, if the probability it is sunny tomorrow is 0.85, then the probability it is not sunny tomorrow is 0.15.

Concept Check ✓ – Answer at <u>cc.dsc10.com</u>


Y= ye = ho

Every time I call my grandma \odot , the probability that she answers her phone is $\frac{1}{3}$, independently for each call. If I call my grandma three times today, what is

the chance that I will talk to her at least once?

- A) 3
- B) $\frac{2}{3}$
- C) $\frac{1}{2}$
- D) [
- E) None of the above.

$$\begin{vmatrix} -\frac{2}{3} \times \frac{2}{3} \\ -\frac{2}{3} \times \frac{2}{3} \end{vmatrix} = \begin{vmatrix} \frac{1}{2} \\ \frac{1}{2} \end{vmatrix}$$



notok


Probability of either of two events happening


ullet Suppose again that A and B are two events, and that all outcomes are equally likely. Then, the probability that either A or B occur is

$$P(A \text{ or } B) = \frac{\# \text{ of outcomes}}{\text{total } \# \text{ of outcomes}} \xrightarrow{\text{satisfying either } A \text{ or } B} \bigvee \bigvee \bigvee$$

• Example 4: I roll a fair six-sided die. What is the probability that the roll/is

even or at least 5?





S happens, then A

The addition rule

- Suppose that if A happens, then B doesn't, and if B happens, then A doesn't.
 - Such events are called mutually exclusive they have no overlap.
- ullet If A and B are any two mutually exclusive events, then

$$P(A \text{ or } B) = P(A) + R(B)$$
 50 Motion

• **Example 5**: Suppose I have two biased coins, a red coin and a blue coin. The red coin flips heads with probability 0.6, and the blue coin flips heads with probability 0.3. I flip both coins once. What's the probability I see two

Aside: Proof of the addition rule for equally-likely events

You are not required to know how to "prove" anything in this course; you may just find this interesting.

If A and B are events consisting of equally likely outcomes, and furthermore A and B are mutually exclusive (meaning they have no overlap), then

$$P(A \text{ or } B) = \frac{\# \text{ of outcomes satisfying either } A \text{ or } B}{\text{total } \# \text{ of outcomes}}$$

$$= \frac{(\# \text{ of outcomes satisfying } A) + (\# \text{ of outcomes satisfying } B)}{\text{total } \# \text{ of outcomes}}$$

$$= \frac{(\# \text{ of outcomes satisfying } A)}{\text{total } \# \text{ of outcomes}} + \frac{(\# \text{ of outcomes satisfying } B)}{\text{total } \# \text{ of outcomes}}$$

$$= P(A) + P(B)$$

Summary, next time

- Probability describes the likelihood of an event occurring.
- There are several rules for computing probabilities. We looked at many special cases that involved equally-likely events.
- There are two general rules to be aware of:
 - The **multiplication rule**, which states that for any two events, $P(A \text{ and } B) = P(B \text{ given } A) \cdot P(A)$.
 - The addition rule, which states that for any two mutually exclusive events, $P(A ext{ or } B) = P(A) + P(B)$.
- Next time: Simulations.